


Seismicity and Earthquakes of Nevada and western U.S.



The 'contemporary' release of accumulated strain registered by Geodesy.

A




With stress – comes strain – and the ultimate and repeated failure of crustal rocks on faults – which in turn produce sound and leave both a geological and geomorphological signature....

B





C




The sudden displacement produces vibrations-sound – which led to development of instruments and arrays to measure those sound to allow determination of the size and location of the source of the earthquakes.

WWSSN Station Map



GSN - Global Seismographic Network



With over 150 permanent station recording continuously this network is aimed at monitoring the largest and longest period seismic waves, and thus the largest earthquakes, around the globe.

The first global seismic network was developed in the 1960's – earthquake vibrations were recorded on paper and collected over months to years and sent to central facility to study particular earthquakes – this has morphed through time to real time recording across a network of more than 150 stations around the world...

- **Partners:**

- [Incorporated Research Institutions for Seismology \(IRIS\)](#)
- [National Science Foundation \(NSF\)](#)
- [University of California San Diego, IDA](#)
- [Comprehensive Test Ban Treaty Organization](#)
- [Federation of Digital Broad-Band Seismograph Networks \(FDSN\)](#)

National Earthquake Information Center (NEIC)

The NEIC collects data through the operation of national and global networks, and through cooperative agreements. To enable the detection and location of all felt earthquakes with the U.S.

the NEIC acts as the National Operations Center of the Advanced National Seismic System (ANSS), a cooperative venture between the NEIC and the operators of the regional seismic networks across the United States.

The NEIC is the national data center and archive for earthquake information.

**QuakeFeed – their app –
should be on your phone**

Locating earthquakes
employs same principles as
locating GPS receivers

Important point is that the
accuracy of

EPICENTER


And

FOCAL DEPTH


Determinations is a function
of how dense is the station
spacing in the networks.

$$\text{Focal depth} = \sqrt{D^2 - \Delta^2}$$


Map View to illustrate epicenter determination



Cross Section to illustrate focal depth determination



This first plot of the global pattern of seismicity by Barazangi and Dorman dates to the 1960s – implication-> the data provide a picture of tectonic strain release at best 'complete' over the last ~50 years....




SEISMICITY OF THE EARTH, 1961-1967, ESSA, CGS EPICENTERS

DEPTHS 000-700 KM.


US Array (reference stations) – A continental-scale network (part of ANSS)


One of 3 components of 'EARTHSCOPE' project funded by NSF – This dates to ~2006...



US Array (Transportable stations)

400 high-quality broadband seismographs leap frogged across the conterminous US.






CISN
California Integrated
Seismic Network

California and western Nevada by the numbers (as of June 2002):


- 614 short-period sensors
- 198 broadband sensors
- 1563 strong-motion sensors (~460 without communications)
- 708 instrumented structures (lifelines, dams, buildings)
- 38 borehole installations




The Southern California Seismic Network (SCSN) records data from more than 370 seismic stations. Each station records seismic waves from both near and distant earthquakes. All the data are transmitted automatically to Caltech/USGS in Pasadena for processing and distribution of information such as epicenters, magnitudes, and ShakeMaps. The SCSN is also part of the California Integrated Seismic Network (CISN) that coordinates earthquake monitoring statewide. The symbols indicate different types of seismic stations.





UNR Seismo Lab 'Nevada Seismic Network'





<http://www.fdsn.org/networks/>






Most earthquakes limited to depths <15-20 km













Orange – broadband
Yellow - broadband
Black – short period
Blue – strong ground motion




University of Utah
Seismograph
Stations.











How often?





Changing topics - Also arising from Seismological Studies – Information of thickness of CRUST



Both little and large earthquakes provide ability to look at velocity structure of the crust.


Depth of MOHO – generally considered the boundary between crust and mantle. It is expressed as a rapid velocity increase and associated with a distinct chemical/mineralogic change.

Map of Crustal Thickness (Depth to Moho). Values range from 30 to 50 km. Blue is thicker, Red is thinner.

Images of crustal variations in the intermountain west

By: Gilbert, HJ; Sheehan, AF

JOURNAL OF GEOPHYSICAL RESEARCH-SOLID EARTH Volume:
109 Issue: B3 Article Number: B03306 Published: MAR 20 2004




Older version of Previous with less resolution but displays broad scale features.

Map of Crustal Thickness (Depth to Moho). Values range from 30 to 50 km. Blue is thicker, Red is thinner.

Images of crustal variations in the intermountain west


By: Gilbert, HJ; Sheehan, AF

JOURNAL OF GEOPHYSICAL RESEARCH-SOLID EARTH Volume:
109 Issue: B3 Article Number: B03306 Published: MAR 20 2004




Geothermal Energy in the USA

GEOHERMAL RESOURCE POTENTIAL




Beneath the Sierra Nevada


Absence of a root in southern



Structure of the Sierra Nevada from receiver functions and implications for lithospheric foundering
 By: Frassetto, Andrew M.; Zandt, George; Gilbert, Hersh; et al.
 GEOSPHERE Volume: 7 Issue: 4 Pages: 898-921 Published: AUG 2011



United States Elevation Map by www.FloodMap.net (beta)



While Crust is 'thin' in Basin and Range – Elevations are high







Figure 4 (a, b) Present-day distribution of gravitational potential energy (ΔPE) in western United States (a). Positive (negative) values of ΔPE indicate lithosphere in a state of extension (compression).

High Elevation leads to increase in vertical stress in B&R- i.e. gravitational potential energy!- and normal faults...





The high topography of the Basin and Range is not smooth – Between the Wasatch and Sierra generally composed of northeast trending ranges and valleys

Valleys on average rest at about 1500m (4500 ft) and highest ranges exceed 3000 m (~10,000 ft)

?s -> when and how did B&R get high and when was topography created

Topography is not same as Relief...

EXPLANATION OF COLORS

- Above 10000 feet
- 8000 to 10000 feet
- 5000 to 8000 feet
- 3000 to 5000 feet
- 1000 to 3000 feet
- 500 to 1000 feet
- Sea Level to 500 feet
- Land Below Sea Level
- To 600 feet deep
- 600 to 4000 feet
- Below 6000 feet

LAND ELEVATIONS AND SEA DEPTHS

POPULATION

- Cities over 1,000,000
- 500,000 to 1,000,000
- 200,000 to 500,000
- 100,000 to 200,000
- Less than 100,000 people

SECTIONAL MAP OF THE UNITED STATES

ABOUT THE MAP:

Scale 25 miles to the inch

Elevations in feet:

- Salt Lakes
- Boundaries
- Marshes
- Rivers
- Canals
- Railroads
- Highways

LOCATOR MAP FOR SIX SECTIONAL MAPS